

# Quantitative Evaluation of NFE2L2/NRF2 and Cytokine Response Induced by Porphyromonas Gingivalis Lipopolysaccharide and Nicotine in a Titanium-Oral Mucosa Model

## Porphyromonas Gingivalis Lipopolisakkarit ve Nikotin Tarafından Uyarılan Titanyum-Ağız Mukoza Modelinde NFE2L2/NRF2 ve Sitokin Yanıtının Kantitatif Olarak Değerlendirilmesi

Gökhan KASNAK<sup>a</sup>, Mervi GÜRSOY<sup>b,c</sup>, Eija KÖNÖNEN<sup>a</sup>, Erhan FIRATLI<sup>d</sup>,  
Stina SYRJÄNEN<sup>b,e</sup>, Ulvi Kahraman GÜRSOY<sup>b</sup>

<sup>a</sup>Istanbul Kent University Faculty of Dentistry, Department of Periodontology, İstanbul, Türkiye

<sup>b</sup>Turku University Institute of Dentistry, Department of Periodontology, Turku, Finland

<sup>c</sup>City of Turku Social Welfare Division, Department of Oral Health, Turku, Finland

<sup>d</sup>Istanbul University Faculty of Dentistry, Department of Periodontology, İstanbul, Türkiye

<sup>e</sup>Turku University Hospital, Clinic of Pathology, Turku, Finland

**ABSTRACT Objective:** Titanium can alter peri-implant mucosa cytokine response to microbiological and chemical stresses. This work used organotypic titanium-oral mucosa model to evaluate oxidant resistance and cytokine response to *Porphyromonas gingivalis* lipopolysaccharide (Pg LPS) and nicotine. **Material and Methods:** Seeding gingival keratinocytes on collagen gels with fibroblasts and adding titanium grade (Ti-Gr) 4, Ti-Gr 5, or hydroxyapatite discs developed an organotypic titanium-oral mucosa model. Pg LPS (1 l/mL), nicotine (1.54 mM), or both were given to the model. The Luminex® xMAPTM technology measured interleukin (IL)-1 $\beta$ , IL-1Ra, IL-8, monocyte chemoattractant protein-1 (MCP-1), and vascular endothelial growth factor levels in culture media. Cultured tissues were paraffin-blocked and immunohistochemistry was used to measure nuclear factor erythroid 2 like 2 (NFE2L2/NRF2), 8-hydroxyguanosine (8-OHdG), and Parkinsonism-associated deglycase (PARK7/DJ-1). Statistical analysis included one-way analysis of variance and Tukey's HSD. **Results:** Nicotine and Pg LPS+nicotine increased IL-1 $\beta$  and IL-1Ra secretions in Ti-Gr 4, Ti-Gr 5, and hydroxyapatite groups (p<0.01). Pg LPS alone (p<0.01) and in conjunction with nicotine (p=0.021) enhanced MCP-1 production, but nicotine alone lowered it (p=0.024). IL-8 concentrations rose in all test groups (p<0.01). All groups treated with nicotine, LPS, or both showed increased 8-Hydroxydeoxyguanosine and NFE2L2/NRF2 immunostainings (p<0.01), whereas PARK7/DJ-1 expression remained unchanged. When assessed without inducers, Ti-Gr 4, Ti-Gr 5, or hydroxyapatite groups showed no significant variations in cytokine secretion patterns, NFE2L2/NRF2, 8-OHdG, or PARK7/DJ-1 immunostainings (p>0.05). **Conclusion:** Titanium did not affect cytokines or oxidant resistance molecules in our organotypic model without inducers. However, Pg LPS, nicotine, and their mixtures enhance cytokines and oxidative resistance molecules.

**Keywords:** Cell culture, three dimensional; keratinocyte; fibroblast; porphyromonas gingivalis lipopolysaccharide; nicotine

**ÖZET Amaç:** Titanyum, peri-implant mukozanın kimyasal ve mikrobiyal uyanlara karşı olan sitokin yanıtını değiştirebilir. Bu çalışmada, organotipik titanyum-ağız mukozası modeli kullanılarak *Porphyromonas gingivalis* lipopolisakkaridi (Pg LPS) ve nikotine karşı oksidatif direnç yanıtının incelenmesi amaçlanmıştır. **Gereç ve Yöntemler:** Organotipik model, fibroblast içeren kolajen jelin üzerine gingival keratinositlerin ekilmesini takiben üzerlerine titanyum grade (Ti-Gr) 4, Ti-Gr 5 ve hidroksiapatit disklerin yerleştirilmesi ile oluşturuldu. Uyarıcı olarak Pg LPS (1 l/mL), nikotin (1,54 mM) veya bileşimleri (Pg LPS+nikotine) modellere uygulandı. Kültür ortamındaki interlökin (IL)-1 $\beta$ , IL-1 reseptör antagonisti (IL-1Ra), IL-8, monosit kemotraktan protein [Monocyte chemoattractant protein-1 (MCP-1)] ve vasküler endotelial büyüme faktörünü seviyeleri Luminex® xMAPTM yöntemi ile ölçüldü. Kültüre edilen örnekler parafin bloklara alınarak, nükleer faktör, eritroid 2 benzeri 2 (NFE2L2/NRF2), 8-hidroksiguanozin (8-OHdG) ve Parkinsonizm ile ilişkilendirilen deglikaz [Parkinsonism-associated deglycase (PARK7/DJ-1)] protein ekspresyonları immünohistokimya ile analiz edildi. İstatistiksel analiz için tek yönlü varyans analizi ve Tukey's HSD testi kullanıldı. **Bulgular:** Ti-Gr 4, Ti-Gr 5 ve hidroksiapatit gruplarında, nikotin ve Pg LPS+nikotin IL-1 $\beta$  ve IL-1Ra ekspresyonunu arttırdığı görüldü (p<0,01). Pg LPS tek başına (p<0,01) ve nikotin ile kombinasyon halinde (p=0,021) uygulandığında MCP-1 seviyesinde artış gözlemlenirken, tek başına nikotin varlığında MCP-1 seviyesinde azalma görüldü (p=0,024). Tüm test gruplarında IL-8 konsantrasyonlarının arttığı tespit edildi (p<0,01). Nikotin, Pg LPS veya bunların kombinasyonunun uygulandığı tüm gruplarda 8-OHdG ve NFE2L2/NRF2 ait immün boyalamalar daha yoğun iken (p<0,01), PARK7/DJ-1 eşit düzeyde gözlemlendi (p>0,05). Herhangi bir uyarıcı olmadığında Ti-Gr 4, Ti-Gr 5 veya hidroksiapatit grupları arasında sitokin seviyeleri ve NFE2L2/NRF2, 8-Hidroksideoksiguanozin veya PARK7/DJ-1 immün boyamalarında herhangi bir fark bulunamadı (p>0,05). **Sonuç:** Organotipik titanyum-ağız mukozası modelinde titanyumun sitokin yanıtı uyarmadığı ve oksidatif direnç moleküllerinin artışına neden olmadığı, ancak Pg LPS, nikotin ve kombinasyonlarının varlığında sitokin yanıtını uyardığı ve oksidatif direnç moleküllerinin ortaya çıkmasına neden olduğu çalışmamızca gösterilmiştir.

**Anahtar Kelimeler:** Hücre kültürü, üç boyutlu; keratinosit; fibroblast; porphyromonas gingivalis lipopolisakkaridler; nikotin

### TO CITE THIS ARTICLE:

Kasnak G, Gürsoy M, Könönen E, Fıratlı E, Syrjänen S, Gürsoy UK. Quantitative evaluation of NFE2L2/NRF2 and cytokine response induced by porphyromonas gingivalis lipopolysaccharide and nicotine in a titanium-oral mucosa model. Türkiye Klinikleri J Dental Sci. 2024;30(4):648-57.

**Correspondence:** Gökhan KASNAK

Istanbul Kent University Faculty of Dentistry, Department of Periodontology, İstanbul, Türkiye

**E-mail:** gkasnak@gmail.com

Peer review under responsibility of Türkiye Klinikleri Journal of Dental Sciences.

**Received:** 26 Mar 2024

**Accepted:** 22 May 2024

**Available online:** 03 Jun 2024

2146-8966 / Copyright © 2024 by Türkiye Klinikleri. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Peri-implant mucositis is a reversible inflammatory condition that affects only the soft tissue around dental implants and, if not treated, might lead to the formation of peri-implantitis.<sup>1</sup> Several risk factors, such as bacterial and chemical insults, take part in the formation of peri-implant mucositis.<sup>2</sup> Bacteria are capable of initiating multiple signaling pathways, including the toll, protein kinase, and complement pathways, which subsequently lead to the synthesis of ROS.<sup>3</sup> Reduced levels of ROS play a crucial role in the intracellular signaling cascade.<sup>4</sup> Conversely, an excessive buildup of ROS produced by pathogenic bacteria leads to oxidative stress, which in turn damages the tissue of the oral mucosa by breaking down its structural components.<sup>3</sup> The biomarker known as 8-Hydroxydeoxyguanosine (8-OHdG) exhibits a high degree of sensitivity in detecting oxidative stress, since it has the ability to accurately indicate even minimal levels of oxidative DNA damage. One potential indication of periodontal tissue breakdown is the observation of an elevated amount of 8-OHdG in both saliva and gingival crevice fluid.<sup>4</sup> In addition to this, the incidence of peri-implant tissue problems is heightened in individuals who smoke, which might also be related to nicotine as an accepted source of exogenous ROS.<sup>1,2,5</sup>

Peri-implant mucosa forms the first barrier against physical, chemical, and microbiological insults, and its integrity is vital for long-term success in implant treatment.<sup>6</sup> Keratinocytes and fibroblasts, which are the cells found in this barrier, regulate the inflammatory response by secreting interleukin-1 (IL-1) and IL-1-receptor antagonist (IL-1Ra). Interleukin-8 (IL-8) and monocyte chemoattractant protein-1 (MCP-1) are also involved in the regulation of neutrophil and macrophage migration.<sup>7</sup> Additionally, they regulate the growth of endothelial cells by secreting vascular endothelial growth factor.<sup>8</sup> Moreover, previous studies have shown evidence that titanium had the ability to alter the immune response in human gingival fibroblasts through its chemical composition and surface properties.<sup>9</sup> There is currently a lack of empirical data about the impact of titanium on the inflammation of the peri-implant mucosa, which is caused by ROS.

The interactions between titanium and oral mucosa cells have been generally studied on monolayer

cell culture models in which cells have been seeded on titanium discs.<sup>10</sup> Although it is possible to obtain consistent and significant results from these in-vitro experiments, they are constrained by the lack of a polarized cell phenotype, insufficient cell differentiation, and limited cell-to-cell interaction.<sup>11</sup> Thus, an organotypic titanium oral mucosa model, which contains both oral keratinocytes and fibroblast cells, can be useful for representing the cellular activities of the peri-implant mucosa.<sup>11</sup> Furthermore, it has been established that three-dimensional cell culture models are preferable to monolayer cell culture models in in-vitro oral research.<sup>12</sup> The main objective of this investigation is to examine the cytokine response, namely IL-1b, IL-1Ra, IL-8, MCP-1, and VEGF, in response to oxidative stress caused by bacterial *Porphyromonas gingivalis* lipopolysaccharide (Pg LPS) and chemical (nicotine) assaults in an organotypic titanium oral mucosa model. Plus, an evaluation was conducted to examine the impact of titanium on the modulation of cytokine activity.


## MATERIAL AND METHODS

### CONSTRUCTION OF THE ORGANOTYPIC TITANIUM ORAL MUCOSA MODEL

The 3D organotypic model resembled the structure of native oral mucosal tissue. It was created by using commercially obtained gingival keratinocytes and fibroblasts, which consisted of three compartments: a collagen matrix at the base, seeded keratinocytes above the base, and the disc placed on top.<sup>12-14</sup> The organotypic oral mucosa-titanium model was constructed in four steps. In the first step, gingival keratinocytes were grown in a medium that had keratinocyte-serum-free medium and antibiotics added (penicillin-streptomycin, 10,000 U/mL, Gibco, Thermo Fisher Sci. Co., Waltham, MA, USA) at 37 °C with a 5% CO<sub>2</sub> concentration. Similarly, gingival fibroblasts were cultured in Dulbecco's modified Eagle's medium (DMEM) at 37 °C and 5% CO<sub>2</sub> concentration. Both cell types were passaged on a weekly basis until they reached a confluence of 80-90%. Secondly, to create the collagen matrix, the gingival fibroblast cells (passages 15 to 20) were suspended in a collagen solution (PureCol, Advanced BioMatrix, Tucson, AZ, USA)

mixed with DMEM at a density of  $3 \times 10^5$ /mL. This suspension was then plated into culture inserts (10 mm diameter, 3 mm pore size, ThinCert, Grenier Bio-One, Monroe, NC, USA). Subsequently, the inserts were meticulously put in 12-well tissue cultivation plates (Corning Inc., Corning, NY, USA) and allowed to harden for a duration of 24 hours. Thirdly, the gingival keratinocytes were cultured on collagen-fibroblast gels at a density of  $8 \times 10^5$ /mL. Finally, when the cells had reached confluence after a seven-day period, inserts were positioned on metal grids to establish the air-liquid interface. After the two days of epithelial cells grown commercially available polished hydroxyapatite (HA, Clarkson Chromatography Inc., South Williamsport, PA, USA), machined surface titanium grade (Ti-Gr) 4 (Cp-Ti, contains 0.4% oxygen, Servo Dental GmbH & Co. KG, Hagen, Germany), and Ti-Gr 5 (Ti-6Al-4V, includes 0.2% oxygen, Implants Dental Implant System Ltd., Trabzon, Türkiye), discs with a height of 2 mm and a diameter of 5 mm were placed on top of the cultures, and the models were allowed to grow for an additional 14 days in antibiotic-free Green's medium for the formation of multi-layered epithelium around the discs. The medium was replaced on a triweekly basis (Figure 1).

In the current study, the experiment involving the cultivation of cell lines did not require any kind of ethical permission because the research was conducted solely through the utilization of cell passages rather than primary cells.



**FIGURE 1:** The illustration of the organotypic 3D Titanium-Oral Mucosa model. After the confluence of epithelial cells grown for two days commercially available titanium and hydroxyapatite disks were placed on the top. After a 14-day incubation period, the models were treated with  $1 \mu\text{L}/\text{mL}$  Porphyromonas gingivalis lipopolysaccharide (Pg LPS), 1.54 mM nicotine (NIC), or their combinations for 24 hours. a) Stimulant (Pg LPS or NIC or Pg LPS+NIC) b) Commercially available titanium/hydroxy apatite disks c) Insert d) 12-well plate e) Medium f) Gingival keratinocyte monolayer g) Collagen layer with gingival fibroblasts h) Air/liquid interface i) Metal grid.

## STIMULATION OF THE ORGANOTYPIC TITANIUM ORAL MUCOSA MODEL AND SAMPLE COLLECTION

At the end of incubation (14-day) of the organotypic oral mucosa Pg LPS ( $1 \mu\text{L}/\text{mL}$ , tlr1-pglps, InvivoGen, San Diego, CA, USA), nicotine (1.54 mM, N0267, Sigma Aldrich, St. Louis, MO, USA) and their combination (Pg LPS+nicotine) were applied to the top of each inserted disc. In control cultures only phosphate-buffered saline was added. The administration quantities of nicotine and Pg LPS were calculated in accordance with the research of Kasnak et al.<sup>15,16</sup> The inserts were extracted from the 12-well plates after a 24-hour incubation period, and the discs were carefully removed. Cytokine analysis was conducted by collecting and storing the medium from each well in a freezer at a temperature of  $-80^\circ\text{C}$ . The oral mucosa samples of organotypic titanium were subjected to fixation in a 10% buffered formalin solution for a duration of 16 hours. Subsequently, the samples were meticulously washed with tap water. Each sample underwent dehydration for 2x30 minutes by gradually increasing the concentration of ethanol. Following a brief xylene cleaning process, all samples were bisected, and the exposed sides of each division were aligned for paraffin embedding.

## ANALYSIS OF CYTOKINE LEVELS

The determination of IL-1 $\beta$ , IL-1Ra, IL-8, MCP-1, and vascular endothelial growth factor (VEGF) levels in the culture media of the organotypic titanium oral mucosa models was conducted using the Luminex<sup>®</sup> xMAP<sup>™</sup> technique (Luminex Corporation, Austin, TX). Commercially accessible kits (pro-human cytokine group I assays; Bio-Rad, Santa Rosa, CA) were employed for this purpose. The assay detects IL-1 $\beta$  at a level of 0.6 pg/mL, IL-1Ra at 5.5 pg/mL, IL-8 at 1.0 pg/mL, MCP-1 at 1.1 pg/mL, and VEGF at 3.1 pg/mL. Just the elute and medium were used to determine the background absorbance. The determination of the detection limit involved identifying the analytical level at which the detected fluorescence intensity signal surpassed the background signal by a margin of two standard deviations.

## IMMUNOHISTOCHEMICAL EXAMINATION OF THE ORGANOTYPIC TITANIUM ORAL MUCOSA MODEL

Immunohistochemistry was performed by cutting sections of paraffin blocks with a thickness of 5  $\mu\text{m}$ . Cytokeratin (M0821, DAKO, Glostrup, Denmark) and vimentin (M7020, DAKO) antibodies were employed for the purpose of localizing epithelial and mesenchymal tissue. To detect ROS, the tissue samples were subjected to immunostaining using 8-OHdG (SC 393870, Santa Cruz Biotechnology, Dallas, TX, USA). Additionally, the anti-oxidative response was identified by immunostaining with nuclear factor erythroid 2 like 2/NFE2L2/NRF2 (SC 722, Santa Cruz Biotechnology, Dallas, TX, USA) and Parkinsonism-associated deglycase (PARK7/DJ-1) (SC 32874, Santa Cruz Biotechnology, Dallas, TX, USA). The usual procedures were employed to perform immunohistochemical stainings for all markers utilizing an automated immunostainer (TechMate, DAKO). Cytokeratin, vimentin, and PARK7/DJ-1 antibodies were diluted at a ratio of 1:50, whilst NFE2L2/NRF2 and 8-OHdG antibodies were diluted at a ratio of 1:75. In summary, the antigen was extracted using a microwave for 5 minutes twice in a citrate buffer with a concentration of 1 mol/L and a pH of 6.0. This was then followed by the inhibition of endogenous peroxidase using 3% H<sub>2</sub>O<sub>2</sub>. The biotinylated secondary antibody (Dako REAL™ Detection System, K5001, DAKO) was used to detect the primary antibody. This antibody was coupled with streptavidin-horseradish peroxidase and visualized using 3,3'-diaminobenzidine tetrahydrochloride in HRP buffer (Dako REAL™ Detection System, K5001, Dako, Glostrup, Denmark).

To assess the immunostainings the regions of interest (ROI) were first determined under microscopy. After this, all ROIs, either left or right contact areas to the discs with unimpaired cell integrity, were scanned using original magnifications ranging from 20X to 100X, and high-resolution pictures were taken. Immunopositivity was graded into three categories: non-stained, moderately stained, and fully stained, based on the intensity of the overall staining for 8-OHdG, PARK7/DJ-1, and NFE2L2/NRF2 to evaluate the ratio of positively stained cells. The validation of the staining intensity of keratinocytes and fibroblasts in


the organotypic oral mucosal model was conducted using the ImageJ software (version 1.46c; Rasband WS, National Institutes of Health, Bethesda, MD, USA) with the immunohistochemical image analysis toolbox plugin version 2 (National Institutes of Health, Bethesda, MD, USA). In a prior investigation, our research team demonstrated the effectiveness of the stains through experimentation on human tissues.<sup>16</sup>

## STATISTICAL ANALYSIS

Statistical analysis was performed by utilizing a statistics software (IBM SPSS Statistics V24.0, IBM Corp, Armonk, New York, USA). The statistical analysis utilized a one-way analysis of variance in conjunction with Tukey's HSD. P-values less than 0.05 were regarded as indicating statistical significance.

## RESULTS

The characterization of the oral mucosa in organotypic titanium model by vimentin, and cytokeratin immunostainings and the surface characteristics were presented as supplementary figures (Figure 1, Figure 2). Induction of oxidative stress by Pg LPS, nicotine, and Pg LPS+nicotine is demonstrated with 8-OHdG immunostainings; the response to those oxidative stimulants is validated with NFE2L2/NRF2 and PARK7/DJ-1 immunostainings. 8-OHdG and NFE2L2/NRF2 stainings were substantially more intense in all disc types after nicotine exposure than in the nontreated controls (8-OHdG stainings  $p=0.028$  for hydroxyapatite,  $p=0.040$  for Ti-Gr 4, and  $p=0.045$  for Ti-Gr 5; NFE2L2/NRF2 stainings  $p=0.033$  for hydroxyapatite,  $p=0.39$  for Ti-Gr 4, and  $p=0.045$  for Ti-Gr 5). Likewise, the combined administration of nicotine and Pg LPS resulted in a significant increase in the 8-OHdG and NFE2L2/NRF2 immunopositivity in all samples derived organotypic titanium-oral mucosa models. The significances of the differences in immunopositivity between exposed and non-exposed samples were as follows: 8-OHdG immunostainings  $p=0.047$  for hydroxyapatite,  $p=0.038$  for Ti-Gr 4, and  $p=0.042$  for Ti-Gr 5; NFE2L2/NRF2 stainings  $p=0.046$  for hydroxyapatite,  $p=0.048$  for Ti-Gr 4, and  $p=0.041$  for Ti-Gr 5). On the other hand, the application only with Pg LPS resulted in an in-



**FIGURE 2:** Images (50X magnification) of titanium-oral mucosal models immunostained with NFE2L2/NRF2, 8-OHdG, and PARK7/DJ-1 antibodies in the absence or presence of irritants (LPS of *Porphyromonas gingivalis*, nicotine, and their combination). 8-OHdG: 8-Hydroxydeoxyguanosine; Ti-Gr: Titanium grade; NFE2L2/NRF2; Nuclear factor erythroid 2 like 2; Pg LPS: *Porphyromonas gingivalis* lipopolysaccharide; PARK7/DJ-1: Parkinsonism-associated deglycase.

crease intensity of 8-OHdG immunostaining in all of the test samples ( $p < 0.01$  for hydroxyapatite,  $p < 0.01$

for Ti-Gr 4, and  $p < 0.01$  for Ti-Gr 5), but no difference was observed in the staining levels of

NFE2L2/NRF2 ( $p>0.05$  for all disc types). Regardless of the stimulant, the PARK7/DJ-1 immunopositivity did not reveal significant differences in any of the specimens (Figure 2 and Figure 3).


According to our cytokine analyses, nicotine significantly increased the levels of IL-1 $\beta$  ( $p<0.01$  for hydroxyapatite, Ti-Gr 4, and Ti-Gr 5), IL-1Ra ( $p<0.01$  for hydroxyapatite, Ti-Gr 4, and Ti-Gr 5), and IL-8 ( $p<0.01$  for hydroxyapatite, Ti-Gr 4, and Ti-Gr 5) in all groups in comparison to their controls. Similarly, the combinational application of Pg LPS and nicotine elevated IL-1 $\beta$  ( $p<0.01$  for hydroxyapatite, Ti-Gr 4, and Ti-Gr 5), IL-1Ra ( $p<0.01$  for hydroxyapatite, Ti-Gr 4, and Ti-Gr 5), IL-8 ( $p<0.01$  for hydroxyapatite, Ti-Gr 4, and Ti-Gr 5) and MCP-1 ( $p=0.021$  for hydroxyapatite,  $p=0.018$  for Ti-Gr 4, and  $p=0.02$  for Ti-Gr 5) levels significantly. Nicotine inhibited the secretion of MCP-1 ( $p=0.024$  for hydroxyapatite,  $p=0.021$  for Ti-Gr 4, and  $p=0.01$  for Ti-Gr 5). At the same time, the Pg LPS increased its secretion in all test groups ( $p<0.01$  for hydroxyapatite, Ti-Gr 4, and Ti-Gr 5). Neither alone nor combinational application of Pg LPS and nicotine did not induce any change in VEGF levels (Figure 4).

Additionally, between-group comparisons were performed based on the disc type, and no difference was observed for any cytokine levels and 8-OHdG, NFE2L2/NRF2, and PARK7/DJ-1 immunostainings (Figure 3, Figure 4).

## DISCUSSION


Soft tissue around dental implants has been studied in different models. Previous research has examined how titanium's surface topography and chemical structure affect peri-implant mucosa. The organotypic titanium oral mucosa model has not been studied for the impact of bacteria and nicotine-induced oxidative stress on cytokine regulation. This study shows significant increases in IL-1Ra, IL-1 $\beta$ , and IL-8 levels after treatment with Pg LPS, nicotine, or both. Nicotine alone decreased MCP-1 levels in the organotypic titanium oral mucosa model, while Pg LPS increased it. We also found that the machined surface Ti-Gr 4 and Ti-Gr 5 modify these cytokines similarly.

The peri-implant soft tissue's principal defense against bacterial and environmental dangers, such as nicotine, is oral keratinocytes and fibroblasts, which respond like healthy gingiva.<sup>6,17</sup> In an organotypic titanium oral mucosa model, gingival keratinocytes and fibroblasts were tested for their cytokine response to Pg LPS and nicotine-induced ROS. Human and animal studies have established the importance of the peri-implant soft tissue.<sup>18,19</sup> However, these study models might have some restrictions due to ethical considerations and complexities in the study protocol. Cell cultures are extensively employed in the evaluation of the biological impact of various substances, but monolayer structure with only one cell



**FIGURE 3:** A comparison of the immunostaining intensities of NFE2L2/NRF2, 8-OHdG, and PARK7/DJ-1 between the control and the organotypic models that were exposed to Pg LPS, NIC, or a combination of the two. The results are given as a bar graph. p values above the bars represents the level of statistical significance in digitally analysed immunopositivity when compared the exposed models with controls.

NFE2L2/NRF2; Nuclear factor erythroid 2 like 2; 8-OHdG:8-Hydroxydeoxyguanosine; PARK7/DJ-1: Parkinsonism-associated deglycase; Pg LPS: Porphyromonas gingivalis lipopolysaccharide; NIC: Nicotine.



**FIGURE 4:** The secreted concentrations of IL-1 $\beta$ , IL-1Ra, IL-8, MCP-1, and VEGF (pg/mL) in the culture media of the organotypic titanium oral mucosa models were illustrated as a bar graph. The significant differences between exposed and nonexposed cultures were indicated with p values above the bars.

IL: Interleukin; IL-1Ra: Interleukin-1receptor antagonist; MCP-1: Monocyte Chemoattractant Protein-1; VEGF: Vascular endothelial growth factor; Pg LPS: Porphyromonas gingivalis lipopolysaccharide.

type do not allow the mimic of the *in-vivo* situation.<sup>20,21</sup> Hence, in order to address these constraints, researchers have employed organotypic oral mucosa models that consist of many layers of epithelial cells.<sup>12</sup> Moreover, the confluence of the epithelial cells, basement membrane, and air are crucial for the modulation of epithelial cell differentiation, which maintains an advantage for the organotypic oral mucosa models over the other *in-vitro* protocols.<sup>22</sup> Based on these facts, Chai WL. et al. have introduced an organotypic oral mucosa model to explore the implant-soft tissue interface.<sup>11,12</sup> The application of Pg LPS, nicotine, and their combination to induce oxidative stress in the organotypic titanium oral mucosa model was a novel approach in this study. On the other hand, the short-term culture protocol, 24 hours, was a study limitation. Despite the inherent limitations of cell culture models, such as the inability to replicate dento-gingival or implant-oral mucosal apparatus, our study may be limited by the fact that we solely applied Pg LPS. LPSs from various periodontopathogens, including the bridging bacteria *Fusobacterium nucleatum*, may potentially have a significant

impact on the development and progression of periodontal and periimplant diseases. Consequently, the building of biofilms or the implementation of LPS mixtures could be helpful for future research.

The role of ROS in periodontal disease has been extensively investigated.<sup>22,23</sup> It has been shown that a rise in ROS levels can be caused by metal ions, which contributes to the activation of fibroblasts and osteoclasts, causes osteolysis surrounding dental implants.<sup>24</sup> Up to date, only in one study, the salivary concentrations of the oxidative biomarkers evaluated in peri-implantitis patients, and it has been reported that total saliva would not be useful to quantify the oxidative damage in peri-implantitis.<sup>25</sup> In the present study, the elevated levels of ROS indicated with the presence of 8-OHdG and the application of nicotine, Pg LPS, and the combination of these factors result in an elevation of IL-1 $\beta$ , IL-8, and IL-1Ra levels. Notably nicotine inhibited the expression of MCP-1, whereas in the presence of Pg LPS, alone or with nicotine, its levels were elevated, and this observation may suggest that the observed effect was contingent upon Pg LPS more than nicotine. Prior

researches have substantiated the presence of heightened levels of IL-8 and MCP-1 subsequent to the incubation of gingival keratinocytes and fibroblasts with *P. gingivalis*.<sup>8,26</sup> Contradictory to that, any change in the IL-8 expression levels in nicotine exposed gingival epithelial cells was not observed by Johnson et al.<sup>27</sup> This might be related to the different methodology; Johnson et al., used gingival epithelial cells as monolayers on organotypic titanium oral mucosa model, which contains both gingival keratinocytes grown on collagen gel containing fibroblasts.<sup>27</sup> Fitzsimmons et al., showed elevated IL-8 and MCP-1 secretions from gingival fibroblasts when incubated with Pg LPS but not that of VEGF which is in line with our results.<sup>8,9</sup>

It is well known that not only *P. gingivalis* is capable of initiating the inflammatory-dependent oxidative response, but also nicotine causes the formation of oxidative stress in several types of oral cells, including gingival keratinocytes and fibroblasts.<sup>28</sup> In the current study, we found that all of the test groups had a significantly higher number of 8-OHdG-stained cells than their respective controls, which is in line with the literature.<sup>29</sup> On the other hand, also the evaluation of the levels of NFE2L2/NRF2, the primary controller of anti-oxidative response, in gingival keratinocytes and fibroblasts might be helpful in understanding the role of the response mechanism against the ROS.<sup>30</sup> To our knowledge, we are the first to study the anti-oxidative response in a titanium-oral mucosa model. Our results showed that the treatment of nicotine and its combination with Pg LPS induced protein expression of NFE2L2/NRF2 in the organotypic model. However, the exposure with Pg LPS only did not stimulate NFE2L2/NRF2 levels. The fact that the antioxidant response against Pg LPS can be controlled by mechanisms other than NFE2L2/NRF2 is one possible explanation for this phenomenon. According to Wang et al. findings, *P. gingivalis* has the ability to activate the forkhead box-O gene, which plays a role in the transcription of antioxidants in gingival epithelial cells.<sup>31</sup> Regarding the levels of PARK7/DJ-1, we did not find any significant differences between the test group and the control group. The in-vitro study may have certain inherent

weaknesses, one of which might be responsible for this outcome.

The cruciality of the interaction between peri-implant mucosa and the dental implant abutment for a long-term successful treatment outcome has been recently reviewed in detail.<sup>32</sup> The adhesion, migration, proliferation, and differentiation of cells (epithelial, fibroblast, osteoblast, and macrophage) may be influenced by changes in the topography, chemistry surface energy, and hydrophilicity of titanium, either individually or in combination. These alterations may occur through modifications in cytokine regulation.<sup>9</sup> Another important finding of the present study was that Ti-Gr 4 (C: 0.08%, N: 0.05%, O: 0.4%, Fe: 0.5%, H: 0.015%, and Ti: balance) and Ti-Gr 5 (C: 0.08%, N: 0.05%, O: 0.2%, Fe: 0.3%, H: 0.015%, Al: 6.75%, V: 4.5%, Y: 0.005%, and Ti: balance) even have different compositions resulted in similar pattern of cytokine release after Pg LPS and nicotine exposures. Furthermore, the effects of Ti-Gr 4 and 5 on cytokine expression in both stimulated and non-stimulated organotypic oral mucosa models are shown to be insignificant when compared to the organotypic oral mucosal model with hydroxyapatite. Our findings suggest that the cytokine expression in the peri-implant mucosa is mostly influenced by *P. gingivalis* and/or nicotine-induced ROS, rather than the type of titanium used. However, due to the complexity of the immune system, even a very minor alteration in the secretion of a cytokine may have biological importance in vivo. The majority of the studies related to cytokine expression mainly focused on the surface properties of the titanium and investigated its effect on different cell types but not gingival keratinocytes. Up to our knowledge, no data compare the modifying effect of Ti-Gr 4 and Ti-Gr 5 on cytokine secretion. Previous studies have shown evidence indicating that titanium elicits the synthesis of pro-inflammatory cytokines, including IL-1b, IL-6, and tumor necrosis factor-alpha (TNF-a), within human macrophages.<sup>33</sup> According to Bruni et al., Ti-Gr 5 stimulated TNF-a secretion in the cultured human umbilical endothelial cells.<sup>34</sup> The primary reason for the discrepancy between our results and those of other researchers may be based on the selection of different cell lines and the design of the cell culture

model. In our study, we examined the impact of titanium on oral gingival keratinocytes and fibroblasts, while Bruni et al. selected umbilical cord vein endothelial cells, and Östberg et al. used neutrophils.<sup>33,34</sup> Changes in surface properties may have been another reason for the inconsistent results. Andrukhov et al.'s study, similar to ours, might provide an answer to this.<sup>35</sup> They compared gingival fibroblasts' cytokine expression between smooth and rough surfaces, as well as between titanium and zirconia surfaces. According to their findings, surface characteristics had only a minor effect on gingival fibroblasts' secretion of IL-8. Thus, newly developed titanium alloys and surface treatments should be tested in further studies.

## CONCLUSION

Exposure to Pg LPS and nicotine increased IL-1 $\beta$ , IL-Ra, and IL-8 secretions by producing ROS in the titanium oral mucosa model. MCP-1 secretion decreased by nicotine but elevated by Pg LPS. Importantly, both Ti-Gr 4 and 5 machined surfaces had similar cytokine secretion patterns.

## Acknowledgments

The authors are grateful to Katja Sampalahti and Oona Hallfors from the Institute of Dentistry, University of Turku, Finland for their outstanding support with cytokine level studies and to Mariia

Valkama from the same institution for her excellent technical assistance in immunohistochemical analyses. We declare that all cell lines used in this study were obtained commercially and were not harvested from any human or animal specifically for this study.

## Source of Finance

This study was supported by the Turku University Foundation (Grant no: 12198, UKG), Finnish Dental Society Apollonia (UKG), and by the Scientific and Technological Research Council of Türkiye (Grant no: BİDEB 2219-1059B191600656, GK).

## Conflict of Interest

No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

## Authorship Contributions

**Idea/Concept:** Gökhan Kasnak, Ulvi Kahraman Gürsoy; **Design:** Gökhan Kasnak, Ulvi Kahraman Gürsoy; **Control/Supervision:** Eija Könönen, Stina Syrjänen, Ulvi Kahraman Gürsoy; **Data Collection and/or Processing:** Gökhan Kasnak, Mervi Gürsoy; **Analysis and/or Interpretation:** Gökhan Kasnak, Mervi Gürsoy, Ulvi Kahraman Gürsoy; **Literature Review:** Gökhan Kasnak; **Writing the Article:** Gökhan Kasnak; **Critical Review:** Mervi Gürsoy, Eija Könönen, Erhan Fıratlı, Stina Syrjänen, Ulvi Kahraman Gürsoy; **References and Fundings:** Gökhan Kasnak, Ulvi Kahraman Gürsoy; **Materials:** Ulvi Kahraman Gürsoy.

## REFERENCES

1. Peri-implant mucositis and peri-implantitis: a current understanding of their diagnoses and clinical implications. *J Periodontol.* 2013;84(4):436-43. [[Crossref](#)] [[PubMed](#)]
2. Jepsen S, Berglundh T, Genco R, Aass AM, Demirel K, Derks J, et al. Primary prevention of peri-implantitis: managing peri-implant mucositis. *J Clin Periodontol.* 2015;42 Suppl 16:S152-7. [[Crossref](#)] [[PubMed](#)]
3. Gözl L, Memmert S, Rath-Deschner B, Jäger A, Appel T, Baumgarten G, et al. Hypoxia and P. gingivalis synergistically induce HIF-1 and NF- $\kappa$ B activation in PDL cells and periodontal diseases. *Mediators Inflamm.* 2015;2015:438085. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
4. Muniz FW, Nogueira SB, Mendes FL, Rösing CK, Moreira MM, de Andrade GM, et al. The impact of antioxidant agents complimentary to periodontal therapy on oxidative stress and periodontal outcomes: A systematic review. *Arch Oral Biol.* 2015;60(9):1203-14. [[Crossref](#)] [[PubMed](#)]
5. Kurgan Ş, Önder C, Altıngöz SM, Bağış N, Uyanık M, Serdar MA, et al. High sensitivity detection of salivary 8-hydroxy deoxyguanosine levels in patients with chronic periodontitis. *J Periodontol Res.* 2015;50(6):766-74. [[Crossref](#)] [[PubMed](#)]
6. Avezov K, Reznick AZ, Aizenbud D. Oxidative damage in keratinocytes exposed to cigarette smoke and aldehydes. *Toxicol In Vitro.* 2014;28(4):485-91. [[Crossref](#)] [[PubMed](#)]
7. Heitz-Mayfield LJ, Lang NP. Comparative biology of chronic and aggressive periodontitis vs. peri-implantitis. *Periodontol 2000.* 2010;53:167-81. [[Crossref](#)] [[PubMed](#)]
8. Basso FG, Soares DG, Pansani TN, Turrioni AP, Scheffel DL, de Souza Costa CA, et al. Effect of LPS treatment on the viability and chemokine synthesis by epithelial cells and gingival fibroblasts. *Arch Oral Biol.* 2015;60(8):1117-21. [[Crossref](#)] [[PubMed](#)]

9. Fitzsimmons TR, Ge S, Bartold PM. Compromised inflammatory cytokine response to *P. gingivalis* LPS by fibroblasts from inflamed human gingiva. *Clin Oral Investig.* 2018;22(2):919-927. [[Crossref](#)] [[PubMed](#)]
10. Schwartz-Filho HO, Morandini AC, Ramos-Junior ES, Jimbo R, Santos CF, Marcantonio E Jr, et al. Titanium surfaces with nanotopography modulate cytokine production in cultured human gingival fibroblasts. *J Biomed Mater Res A.* 2012;100(10):2629-36. [[Crossref](#)] [[PubMed](#)]
11. Lagneau C, Farges JC, Exbrayat P, Lissac M. Cytokeratin expression in human oral gingival epithelial cells: in vitro regulation by titanium-based implant materials. *Biomaterials.* 1998;19(11-12):1109-15. [[Crossref](#)] [[PubMed](#)]
12. Chai WL, Moharamzadeh K, Brook IM, Emanuelsson L, Palmquist A, van Noort R. Development of a novel model for the investigation of implant-soft tissue interface. *J Periodontol.* 2010;81(8):1187-95. [[Crossref](#)] [[PubMed](#)]
13. Gursoy UK, Pöllänen M, Könönen E, Uitto VJ. A novel organotypic dento-epithelial culture model: effect of *Fusobacterium nucleatum* biofilm on B-defensin-2, -3, and LL-37 expression. *J Periodontol.* 2012;83(2):242-7. [[Crossref](#)] [[PubMed](#)]
14. Mäkelä M, Larjava H, Pirilä E, Maisi P, Salo T, Sorsa T, et al. Matrix metalloproteinase 2 (gelatinase A) is related to migration of keratinocytes. *Exp Cell Res.* 1999;251(1):67-78. [[Crossref](#)] [[PubMed](#)]
15. Oksanen J, Hormia M. An organotypic in vitro model that mimics the dento-epithelial junction. *J Periodontol.* 2002;73(1):86-93. [[Crossref](#)] [[PubMed](#)]
16. Kasnak G, Firatli E, Könönen E, Olgac V, Zeidán-Chuliá F, Gursoy UK. Elevated levels of 8-OHdG and PARK7/DJ-1 in peri-implantitis mucosa. *Clin Implant Dent Relat Res.* 2018;20(4):574-82. [[Crossref](#)] [[PubMed](#)]
17. Berglundh T, Lindhe J, Marinello C, Ericsson I, Liljenberg B. Soft tissue reaction to de novo plaque formation on implants and teeth. An experimental study in the dog. *Clin Oral Implants Res.* 1992;3(1):1-8. [[Crossref](#)] [[PubMed](#)]
18. Schupbach P, Glauser R. The defense architecture of the human peri-implant mucosa: a histological study. *J Prosthet Dent.* 2007;97(6 Suppl):S15-25. Erratum in: *J Prosthet Dent.* 2008;99(3):167. [[Crossref](#)] [[PubMed](#)]
19. Hermann JS, Buser D, Schenk RK, Schoolfield JD, Cochran DL. Biologic Width around one- and two-piece titanium implants. *Clin Oral Implants Res.* 2001;12(6):559-71. [[Crossref](#)] [[PubMed](#)]
20. Groeger SE, Meyle J. Epithelial barrier and oral bacterial infection. *Periodontol 2000.* 2015;69(1):46-67. [[Crossref](#)] [[PubMed](#)]
21. Imamura K, Kokubu E, Kita D, Ota K, Ishihara K, Saito A. Cigarette smoke condensate modulates migration of human gingival epithelial cells and their interactions with *Porphyromonas gingivalis*. *J Periodontol Res.* 2015;50(3):411-21. [[Crossref](#)] [[PubMed](#)]
22. Chapple IL, Matthews JB. The role of reactive oxygen and antioxidant species in periodontal tissue destruction. *Periodontol 2000.* 2007;43:160-232. [[Crossref](#)] [[PubMed](#)]
23. Meyle J, Chapple I. Molecular aspects of the pathogenesis of periodontitis. *Periodontol 2000.* 2015;69(1):7-17. [[Crossref](#)] [[PubMed](#)]
24. Żukowski P, Maciejczyk M, Waszkiel D. Sources of free radicals and oxidative stress in the oral cavity. *Arch Oral Biol.* 2018;92:8-17. [[Crossref](#)] [[PubMed](#)]
25. Sánchez-Siles M, Lucas-Azorin J, Salazar-Sánchez N, Carbonell-Meseguer L, Camacho-Alonso F. Salivary Concentration of Oxidative Stress Biomarkers in a Group of Patients with Peri-Implantitis: A Transversal Study. *Clin Implant Dent Relat Res.* 2016;18(5):1015-22. [[Crossref](#)] [[PubMed](#)]
26. Kusumoto Y, Hirano H, Saitoh K, Yamada S, Takedachi M, Nozaki T, et al. Human gingival epithelial cells produce chemotactic factors interleukin-8 and monocyte chemoattractant protein-1 after stimulation with *Porphyromonas gingivalis* via toll-like receptor 2. *J Periodontol.* 2004;75(3):370-9. [[Crossref](#)] [[PubMed](#)]
27. Johnson GK, Guthmiller JM, Joly S, Organ CC, Dawson DV. Interleukin-1 and interleukin-8 in nicotine- and lipopolysaccharide-exposed gingival keratinocyte cultures. *J Periodontol Res.* 2010;45(4):583-8. [[Crossref](#)] [[PubMed](#)]
28. Nguyen TT, Huynh NN, Seubbuk S, Nilmoje T, Wanasuntronwong A, Surarit R. Oxidative stress induced by *Porphyromonas gingivalis* lysate and nicotine in human periodontal ligament fibroblasts. *Odontology.* 2019;107(2):133-41. [[Crossref](#)] [[PubMed](#)]
29. Dionigi C, Larsson L, Carcuac O, Berglundh T. Cellular expression of DNA damage/repair and reactive oxygen/nitrogen species in human periodontitis and peri-implantitis lesions. *J Clin Periodontol.* 2020;47(12):1466-75. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
30. Cao H, Wang L, Chen B, Zheng P, He Y, Ding Y, et al. DNA demethylation upregulated Nrf2 expression in alzheimer's disease cellular model. *Front Aging Neurosci.* 2016;7:244. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
31. Wang Y, Zhou Y, Graves DT. FOXO transcription factors: their clinical significance and regulation. *Biomed Res Int.* 2014;2014:925350. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
32. Guo T, Gulati K, Arora H, Han P, Fournier B, Ivanovski S. Orchestrating soft tissue integration at the transmucosal region of titanium implants. *Acta Biomater.* 2021;124:33-49. [[Crossref](#)] [[PubMed](#)]
33. Östberg AK, Dahlgren U, Sul YT, Johansson CB. Inflammatory cytokine release is affected by surface morphology and chemistry of titanium implants. *J Mater Sci Mater Med.* 2015;26(4):155. [[Crossref](#)] [[PubMed](#)]
34. Bruni S, Martinesi M, Stio M, Treves C, Bacci T, Borgioli F. Effects of surface treatment of Ti-6Al-4V titanium alloy on biocompatibility in cultured human umbilical vein endothelial cells. *Acta Biomater.* 2005;1(2):223-34. [[Crossref](#)] [[PubMed](#)]
35. Andrukhov O, Behm C, Blufstein A, Wehner C, Gahn J, Pippenger B, et al. Effect of implant surface material and roughness to the susceptibility of primary gingival fibroblasts to inflammatory stimuli. *Dent Mater.* 2020;36(6):e194-e205. [[Crossref](#)] [[PubMed](#)]